

Patient Reminder Checklist



No intercourse for 48 hours before exam.



No vaginal medication, personal lubricants or douching 48 hours before exam.



Schedule appointment 2 weeks after first day of the patient's last menstrual period.

Scan below to access Pap test collection videos and complete quick reference guides for each device.




Hologic provides this collection procedure guide as a general informational tool only; it is not an affirmative instruction or guarantee of performance. It is the sole responsibility of the clinician to read and understand the appropriate package insert and comply with applicable local, state and federal rules and regulations.

References **1.** Cervicovaginal Cytology Based on the Papanicolaou Technique; Approved Guideline – Third Edition (Clinical and Laboratory Standards Institute GP15-A3). **2.** Hologic internal study,data on file. **3.** Rovers Cervex-Brush Combi, Information For Use, 2010-11.

hologic.com | diagnostic.solutions@hologic.com | +1.888.484.4747

PB-00803-001 Rev. 003 © 2024 Hologic, Inc. All rights reserved. Hologic, PreservCyt, ThinPrep and associated logos are trademarks and/or registered trademarks of Hologic, Inc. and/or its subsidiaries in the United States and/or other countries. All other trademarks are the property of their respective owners. This information is intended for medical professionals in the U.S. and other markets and is not intended as a product solicitation or promotion where such activities are prohibited. Because Hologic materials are distributed through websites, eBroadcasts and tradeshows, it is not always possible to control where such materials appear. For specific information on what products are available for sale in a particular country, please contact your Hologic representative or write to diagnostic.solutions@hologic.com.

ThinPrep®
PAP TEST




ThinPrep®
PAP TEST

A Guide to Success with the ThinPrep® Pap Test by Chosen Device

1 Prepare Speculum^{1,2}

► Warm water is preferred



► Only use carbomer-free lubricants

If lubricant is desired, the following lubricants are validated for use with the ThinPrep® Pap Test.

Lubricant	Manufacturer
✓ Pap Test Lubricating Jelly	Aseptic Control Products
✓ SurgiLube Surgical Lubricant	HR Pharmaceuticals
✓ CerviLube	Sion Biotech

Contact your medical supplies distributor for purchasing details.

► Dime-sized amount

3 Rinse the Chosen Device Immediately

► Sanitary collection - fingers remain on the handle, not the brush




Push the Broom into the bottom of the vial 10 times, forcing the bristles apart.



Swirl the Spatula vigorously into the bottom of the vial 10 times.

The ThinPrep® collection process preserves and fixes the cells in the solution, not the device head.



Rotate the Brush in the solution 10 times while pushing it against the PreservCyt® vial wall.


2 Obtain Sample

Sample collection for the ThinPrep® Pap Test may be performed with a Broom-like, Spatula/Brush, or Rovers® Cervex-Brush® Combi device.

► Rotate per manufacturer's guidance

► Do not over rotate


A Clinician Collection with the Broom-like Device




Rotate clockwise
360° 5x

Push gently and rotate the broom in a clockwise direction for five complete, 360 degree turns.

B Clinician Collection with the Spatula/Brush Device




Rotate the contoured end
360° 1x



Rotate in one direction
1/4 to 1/2

Insert the brush into the cervix until only the bottom-most fibers are exposed. Slowly rotate 1/4 or 1/2 turn in one direction.


C Clinician Collection with the Rovers® Cervex-Brush® Combi Device




Rotate clockwise
360° 2x

Push gently, and rotate the brush in a clockwise direction 2 times for 2 complete 360° degree turns.³


4 Remove and Discard the Collection Device



Do not pop off the head of the broom.



Do not clip the head of the brush.



Do not pop off the head of the Rovers.
Do not clip the head of the Rovers.

No collection device should be left in the vial.